arXiv:1502.04443v1 [quant-ph] 16 Feb 2015

Entanglement of simultaneous and non-simultaneous Accelerated Qubit-Qutrit systems

N. Metwally
[1] Department of Mathematics, College of Science, Bahrain University, Bahrain
[2]Department of Mathematics, Faculty of Science, Aswan University, Aswan, Egypt
email:nmetwally@gmail.com, nmetwally@uob.edu.bh
Tel:00973 33237474, Fax: 00973 17449145

Abstract

In this contribution, we investigate the entanglement behavior of a composite system consists of two different dimensional subsystems in non-inertial frames. In particular, we consider a composite system of qubit(two-dimensional) subsystem, and qutrit, (three-dimensional) subsystem. The degree of entanglement is quantified for different cases, where it is assumed that the two-subsystems are simultaneously or non- simultaneously accelerated. The entanglement decays as the acceleration of any subsystem increases.In general, the decay rate of entanglement increases as the dimensional of the accelerated subsystem increases. These results could be important in building an accelerated quantum network consists of different dimensions nodes.

Keyword: non-inertial frames, Entanglement, acceleration.

1 Introduction

It is well known that, quantum correlations are the most important physical quantities that have many applications in the context of quantum information, communication and computations [1, 2, 3]. The behavior of entanglement between different systems has investigated for small and high dimensions systems [4]. However, the properties of systems which are consists of different dimensional subsystems have investigated in different ways. For example, the behavior of entanglement for a two-parameter class of qubit-qutrit system passes through dissipative channels has discussed by Wei et.al, [5]. Karat and Gedik [6] have discussed the quantum and classical correlation of qubit-qutrit systems in the presence of classical dephasing environment. The quantum correlation's behavior of a qubit-qutrit system under the effect of dephasing and bit flip channels has investigated by Guo et. al., [7]. Moreover, X. Xiao [8] has investigated the possibility of protecting the quantum correlation of qubit-qutrit system by using weak measurement and reversal.

Recently, the behavior of entanglement in non-inertial frames has investigated for different dimensional systems. For example, Alsing et. al., [14] have discussed the decay of entanglement between two modes of a free Dirac field in a non-inertial frame. The correlation between two modes of Dirac fields in non-inertial frames have investigated by Wang et. al, [10]. The dynamics of a general two qubit system in non-inertial frame is investigated by Metwally [11]. The possibility of using an accelerated class of X-state to perform quantum coding is discussed in [12]. Moreover, teleportation in non-inertial frame is discussed by several authors see for example ([13, 14, 15]).

However, for composite system which consists of different dimensions subsystems, there are some limited efforts have been done to study the entangled properties of these systems in non-inertial frames. The effect of global, collective, local and multilocal depolarizing noise on qutrit-qutrit states is studied by S. Khan and M. Khan [16]. The measurement induced disturbance for qubit-qutrit system in non-inertial frames under the effect of different noise that a source supplies randomly some users who located in different nodes with different dimensional states to generated entangled network [18]. So, may any two nodes of the same or different dimensions systems are connected. Then it is important to quantify the degree of entanglement to find the optimal connection. Therefor, we are motivated to consider a general class of qubit-qutrit system in non-inertial frame, where we assume that these subsystems simultaneous and non-simultaneous accelerated.

This manuscript is organized as follows: In Sec. 2, we define the general form of the suggested system. The final state of the system is obtained analytically when one or both subsystems are accelerated simultaneously or non- simultaneously. In Sec. 3, the survival of entanglement between the two subsystems is quantified by using the negativity as a measure of entanglement. Finally, the results are summarized in Sec. 4.

2 System and its evolution

In this contribution, it is assumed that we have a composite system, consists of qubit (A) and qutrit (B), is given in its general form. The evolution the final state of the qubit-qutrit (Q_{bt}) system is investigated for three different cases: (i) the subsystem (A) (qubit) is accelerated while qutrit (B) is in a rest,(ii) the subsystem (B) (qutrit) is accelerated while the subsystem (A) is in a rest, (iii) Both subsystems are accelerated. The general state of the qubit-qutrit system is given by,

$$\rho_{Q_{bt}} = \frac{1}{6}\Big\{I_2 \otimes I_3 + \sum_{i=1}^{3} s_i\sigma_i \otimes I_3 + \sum_{i=1}^{8} t_j\tau_j \otimes I_3 + \sum_{i=1}^{3}\sum_{j=1}^{8} c_{ij}\sigma_i \otimes \tau_j\Big\}, \tag{1}$$

where I_2 and I_3 are the unit matrix of size 2×2 and 3×3 respectively, $s_i = tr\{\rho_{Q_{bt}}\sigma_i \otimes I_3\}$, $t_j = tr\{\rho_{Q_{bt}} I_2 \otimes \tau_j\}$ are the Bloch vectors for the qubit and the qutrit respectively, $i = 1..3$, $j = 1..8$. The operators σ_i and τ_j are the traceless hermitian generators of $Su(2)$ and $Su(3)$ respectively. The elements $c_{ij} = tr\{\rho_{Q_{bt}}\sigma_i \otimes \tau_j\}$, represent the correlation matrix between the qubit (A) and the qutrit (B). In the computational basis 0, 1 and 2, the operators σ_i and τ_j can be written as,

$$\begin{aligned}
\sigma_1 &= |0\rangle\langle 1| + |1\rangle\langle 0|, \quad \sigma_2 = i|0\rangle\langle 1| - |1\rangle\langle 0|), \quad \sigma_3 = |1\rangle\langle 1| - |0\rangle\langle 0|, \\
\tau_1 &= |0\rangle\langle 1| + |1\rangle\langle 0|, \quad \tau_2 = i(|1\rangle\langle 0| - |0\rangle\langle 1|), \quad \tau_3 = |0\rangle\langle 0| - |1\rangle\langle 1|, \\
\tau_4 &= |0\rangle\langle 2| + |2\rangle\langle 0|, \quad \tau_5 = i(|2\rangle\langle 0| - \langle 0||2\rangle), \quad \tau_6 = |1\rangle\langle 2| + |2\rangle\langle 1|, \\
\tau_7 &= i(|2\rangle\langle 1| - |1\rangle\langle 2|), \quad \tau_8 = (|0\rangle\langle 0| + |1\rangle\langle 1| - 2|2\rangle\langle 2|)/\sqrt{3}.
\end{aligned} \tag{2}$$

In the computational basis, $|00\rangle, |01\rangle, |02\rangle, |10\rangle, |11\rangle$ and $|12\rangle$, the state (1) can be written as,

$$\begin{aligned}
\varrho_{00,ij} &= \mathcal{A}_{\ell_1}, \quad \ell_1 = 1...6, \qquad \varrho_{01,ij} = \mathcal{A}_{\ell_2}, \quad \ell_2 = 7...12, \\
\varrho_{02,ij} &= \mathcal{A}_{\ell_3} \quad \ell_3 = 13...18, \quad \varrho_{10,ij} = \mathcal{A}_{\ell_4}, \quad \ell_4 = 19...24, \\
\varrho_{11,ij} &= \mathcal{A}_{\ell_5 1}, \quad \ell_5 = 25...30, \quad \varrho_{12,ij} = \mathcal{A}_{\ell_6}, \quad \ell_6 = 31...36,
\end{aligned} \tag{3}$$

where $ij = 00, 01, 02, 10, 11, 12$ and $\mathcal{A}_\ell, \ell = 1...36$ are given in the appendix (A). For example, the element $\varrho_{00,00} = \mathcal{A}_1$, $\varrho_{00,01} = \mathcal{A}_2$, $\varrho_{02,02} = \mathcal{A}_{15}$ $\varrho_{12,12} = \mathcal{A}_{36}$ and so on.

Now we investigate the evolution of the Q_{bt} state (1) in non-inertial frame when one or both of its subsystems are accelerated as following:

1. *Only the qubit is accelertaed*

   In this context, it is important to review the behavior of a particle in non-inertial frames. It has been shown that, in the inertial frames, Minkowsik coordinates (t, z) are used to describe Dirac field, while in the uniformly accelerated case, Rindler coordinates (τ, x) are more adequate. The relations between the Minkowski and Rindler coordinates are given by[13, 19],

$$\tau = r\ tanh\left(\frac{t}{z}\right), \quad x = \sqrt{t^2 - z^2}, \tag{4}$$

   where $-\infty < \tau < \infty$, $-\infty < x < \infty$ and r is the acceleration of the moving particle. The relations (4) describe two regions in Rindler's spaces: the first region I for $|t| < x$ and the second region II for $x < -|t|$.

   A single mode k of fermions and anti-fermions in Minkowski space is described by the annihilation operators a_{kU} and b_{-kU} respectively. In terms of Rindler's operators ($c^{(I)}_{kR}, d^{(II)}_{-kL}$), the Minkowski operators can be written as [20, 21],

$$\begin{aligned} a_{kU} &= \cos r c^{(I)}_{k,R} - \exp(-i\phi)\sin r d^{(II)}_{-k,L}, \\ b^{\dagger}_{-kU} &= \exp(i\phi)\sin r c^{(I)}_{k,R} + \cos r d^{(II)}_{k,L}, \end{aligned} \tag{5}$$

   where, $tan\ r = e^{-\pi\omega\frac{c}{a}}$, $0 \leq r \leq \pi/4$, a is the acceleration such that $0 \leq a \leq \infty$, ω is the frequency of the travelling qubits, c is the speed of light, and ϕ is an unimportant phase that can be absorbed into the definition of the operators [22]. The operators (5) mix a particle in region I and an anti particle in region II as,

$$\begin{aligned} |0_k\rangle &= \cos r|0_k\rangle_I|0_{-k}\rangle_{II} + \sin r|1_k\rangle_I|1_{-k}\rangle_{II}, \\ |1_k\rangle &= a^{\dagger}_k|0_k\rangle = |1_k\rangle_I|0_k\rangle_{II}. \end{aligned} \tag{6}$$

   Now, by using the initial state of the qubit-qutrit system $\rho_{Q_{bt}}$ (1) and the operators (6), the final state in the first region I of the Rindler space can be written in the basis $00, 01, 02, 10, 11$ and 12. The density operator is represented by a matrix of size 6×6. Its elements are given by,

$$\begin{aligned}
\rho^{(q)}_{00,00} &= c_q^2\varrho_{00,00}, \quad \rho^{(q)}_{00,01} = c_q^2\varrho_{00,01}, \quad \rho^{(q)}_{00,02} = c_q^2\varrho_{00,02}, \quad \rho^{(q)}_{00,10} = c_q\varrho_{00,10}, \\
\rho^{(q)}_{00,11} &= c_q\varrho_{00,11}, \quad \rho^{(q)}_{00,12} = c_q\varrho_{00,12}, \quad \rho^{(q)}_{01,00} = c_q^2\varrho_{01,00}, \quad \rho^{(q)}_{01,01} = c_q^2\varrho_{01,01}, \\
\rho^{(q)}_{01,02} &= c_q^2\varrho_{01,02}, \quad \rho^{(q)}_{01,10} = c_q\varrho_{01,10}, \quad \rho^{(q)}_{01,11} = c_q\varrho_{01,11}, \quad \rho^{(q)}_{01,12} = c_q\varrho_{01,12}, \\
\rho^{(q)}_{02,00} &= c_q^2\varrho_{02,00}, \quad \rho^{(q)}_{02,01} = c_q^2\varrho_{02,01}, \quad \rho^{(q)}_{02,02} = c_q^2\varrho_{02,02}, \quad \rho^{(q)}_{02,10} = c_q\varrho_{02,10}, \\
\rho^{(q)}_{02,11} &= c_q\varrho_{02,11}, \quad \rho^{(q)}_{02,12} = c_q\varrho_{02,12}, \quad \rho^{(q)}_{10,00} = c_q\varrho_{10,00}, \quad \rho^{(q)}_{10,01} = c_q\varrho_{10,01}, \\
\rho^{(q)}_{10,02} &= c_q\varrho_{10,02}, \quad \rho^{(q)}_{10,10} = s_q^2\varrho_{00,00} + \varrho_{10,10}, \quad \rho^{(q)}_{10,11} = s_q^2\varrho_{00,01} + \varrho_{10,11}, \\
\rho^{(q)}_{10,12} &= s_q^2\varrho_{00,02} + \varrho_{10,12}, \quad \rho^{(q)}_{11,00} = c_q\varrho_{11,00}, \quad \rho^{(q)}_{11,01} = c_q\varrho_{11,01}, \quad \rho^{(q)}_{11,02} = c_q\varrho_{11,02}, \\
\rho^{(q)}_{11,10} &= s_q^2\varrho_{01,00} + \varrho_{11,10}, \quad \rho^{(q)}_{11,11} = s_q^2\varrho_{01,01} + \varrho_{11,11}, \quad \rho^{(q)}_{11,12} = s_q^2\varrho_{01,02} + \varrho_{11,12}, \\
\rho^{(q)}_{12,00} &= c_q\varrho_{12,00}, \quad \rho^{(q)}_{12,01} = c_q\varrho_{12,01}, \quad \rho^{(q)}_{12,02} = c_q\varrho_{12,02}, \quad \rho^{(q)}_{12,10} = s_q^2\varrho_{02,00} + \varrho_{12,10}, \\
\rho^{(q)}_{12,11} &= s_q^2\varrho_{02,01} + \varrho_{12,11}, \quad \rho^{(q)}_{12,12} = s_q^2\varrho_{02,02} + \varrho_{12,12},
\end{aligned} \tag{7}$$

   where $c_q = \cos r_q$ and $s_q = sinr_q$.

2. *Only the qutrit is accelerated*
   In this case, it is assumed that the qubit is in an inertial frame, while the qutrit undergoes a constant acceleration. For the two level qubit system (spin 1/2) particles, the Pauli-principle allows only two occupation numbers i.e.,”0” and ”1” [14]. However, the possibility of allowing the occupation number ”2” is discussed by Leòn and Martìnez [23]. Under the single mode-approximation, Minlowaski vacuum state $|0_M\rangle$ in the Rindler space reads as,

$$|0_M\rangle = cos^2 r|0\rangle_I|0\rangle_{II} + e^{i\phi}\sin r\cos r(|\mathcal{U}\rangle_I|\mathcal{D}\rangle_{II} + |\mathcal{D}\rangle_I|\mathcal{U}\rangle_{II}) + e^{2i\phi}sin^2 r|\mathcal{D}\rangle_I|\mathcal{P}\rangle_{II}, \quad (8)$$

   where $|\mathcal{U}\rangle, |\mathcal{D}\rangle$ and $|\mathcal{P}\rangle$ are the spin up, spin down and pair state respectively. The spin up and down in Rindler space are given by,

$$\begin{aligned} |\mathcal{U}_M\rangle &= cosr|\mathcal{U}\rangle_I|0\rangle_{II} + e^{i\phi}\sin r|\mathcal{P}\rangle_I|\mathcal{U}\rangle_{II}, \\ |\mathcal{D}_M\rangle &= cosr|\mathcal{D}\rangle_I|0\rangle_{II} - e^{i\phi}\sin r|\mathcal{P}\rangle_I|\mathcal{D}\rangle_{II}. \end{aligned} \quad (9)$$

   By using the initial state (1) and the transformation (8,9) one can obtain the final state of the accelerated system. After tracing out the mode in the second region II, we obtain the final accelerated state in the first region I of the Rindler space. In the basis, $|\ell, 0\rangle, |\ell, \mathcal{D}\rangle, |\ell, \mathcal{U}\rangle, |\ell, \mathcal{P}\rangle$ $(\ell = 0, 1)$, the density operator is defined by a matrix of size 8×8, where its elements are given by,

$$\begin{aligned}
\rho^{(t)}_{00,00} &= c_t^4\varrho_{00,00}, \quad \rho(t)_{00,0\mathcal{D}} = c_t^3\varrho_{00,01}, \quad \rho(t)_{00,0\mathcal{U}} = c_t^2\varrho_{00,02}, \quad \rho(t)_{00,0\mathcal{P}} = 0, \\
\rho(t)_{00,10} &= c_t^4\varrho_{00,10}, \quad \rho^{(t)}_{00,1\mathcal{D}} = c_t^3\varrho_{00,11}, \quad \rho^{(t)}_{00,1\mathcal{U}} = c_t^2\varrho_{00,12}, \quad \rho_{00,1\mathcal{P}} = 0, \quad \rho^{(t)}_{0\mathcal{D},00} = c_t^4\varrho_{01,00}, \\
\rho^{(t)}_{0\mathcal{D},0\mathcal{D}} &= c_t^2(s_t^2\varrho_{00,00} + \varrho_{01,01}), \quad \rho^{(t)}_{0\mathcal{D},0\mathcal{U}} = c_t^2\varrho_{01,02}, \quad \rho_{0\mathcal{D},0\mathcal{P}} = c_t s_t^2\varrho_{00,02}, \\
\rho^{(t)}_{0\mathcal{D},\mathcal{D}0} &= c_t^4\varrho_{01,10}, \quad \rho^{(t)}_{0\mathcal{D},\mathcal{D}\mathcal{D}} = c_t^2(s_t^2\varrho_{00,10} + \varrho_{01,11}), \quad \rho^{(t)}_{0\mathcal{D},1\mathcal{U}} = c_t^2\varrho_{01,12}, \quad \rho^{(t)}_{0\mathcal{D},1\mathcal{P}} = c_t s_t^2\varrho_{00,12}, \\
\rho^{(t)}_{0\mathcal{U},00} &= c_t^3\varrho_{02,00}, \quad \rho^{(t)}_{02\mathcal{U},0\mathcal{D}} = c_t^3\varrho_{02,01}, \quad \rho^{(t)}_{0\mathcal{U},0\mathcal{U}} = c_t^2(s_t^2\varrho_{00,01} + \varrho_{02,02}), \quad \rho^{(t)}_{0\mathcal{U},0\mathcal{P}} = -c_t s_t^2\varrho_{02,10}, \\
\rho^{(t)}_{0\mathcal{U},10} &= c_t^3\varrho_{02,10}, \quad \rho^{(t)}_{0\mathcal{U},1\mathcal{D}} = c_t^2\varrho_{02,11}, \quad \rho^{(t)}_{0\mathcal{U},1\mathcal{U}} = c_t^2(s_t^2\varrho_{02,10} + \varrho_{02,12}), \quad \rho^{(t)}_{0\mathcal{U},1\mathcal{P}} = -c_t s_t^2\varrho_{00,11}, \\
\rho^{(t)}_{0\mathcal{P},00} &= 0, \quad \rho^{(t)}_{0\mathcal{P},0\mathcal{D}} = c_t s_t^2\varrho_{02,00}, \quad \rho^{(t)}_{0\mathcal{P},0\mathcal{U}} = -c_t s_t^2\varrho_{01,00}, \quad \rho^{(t)}_{0\mathcal{P},0\mathcal{P}} = s_t^2(s_t^2\varrho_{00,00} + \varrho_{01,02}), \\
\rho^{(t)}_{0\mathcal{P},10} &= 0, \quad \rho^{(t)}_{0\mathcal{P},1\mathcal{D}} = c_t s_t^2\varrho_{02,10}, \quad \rho^{(t)}_{0\mathcal{P},1\mathcal{U}} = -c_t s_t^2\varrho_{01,10}, \quad \rho^{(t)}_{0\mathcal{P},1\mathcal{P}} = s_t^4\varrho_{00,10}, \\
\rho^{(t)}_{10,00} &= c_t^4\varrho_{10,00}, \quad \rho^{(t)}_{10,0\mathcal{D}} = c_t^3\varrho_{10,01}, \quad \rho^{(t)}_{10,0\mathcal{U}} = c_t^2\varrho_{10,02}, \quad \rho^{(t)}_{10,0p} = 0 \\
\rho^{(t)}_{10,10} &= c_t^4\varrho_{10,10}, \quad \rho^{(t)}_{10,1\mathcal{D}} = c_t^3\varrho_{10,11}, \quad \rho^{(t)}_{10,12\mathcal{U}} = c_t^2\varrho_{10,12}, \quad \rho^{(t)}_{10,1\mathcal{P}} = 0, \\
\rho^{(t)}_{1\mathcal{D},00} &= c_t^3\varrho_{11,00}, \quad \rho^{(t)}_{1\mathcal{D},0\mathcal{D}} = c_t^2\varrho_{11,01}, \quad \rho^{(t)}_{1\mathcal{D},0\mathcal{U}} = c_t^2\varrho_{11,02}, \quad \rho^{(t)}_{1\mathcal{D},0\mathcal{P}} = c_t s_t^2\varrho_{10,02}, \\
\rho^{(t)}_{1\mathcal{D},10} &= c_t^3\varrho_{11,10}, \quad \rho^{(t)}_{1\mathcal{D},1\mathcal{D}} = c_t^2 s_t^2\varrho_{11,11}, \quad \rho^{(t)}_{1\mathcal{D},1\mathcal{U}} = c_t^2\varrho_{11,12}, \quad \rho^{(t)}_{1\mathcal{D},1\mathcal{P}} = c_t^2 s_t^2\varrho_{10,12}, \\
\rho^{(t)}_{1\mathcal{U},00} &= c_t^2\varrho_{12,00}, \quad \rho^{(t)}_{1\mathcal{U},0\mathcal{D}} = c_t^2\varrho_{12,01}, \quad \rho^{(t)}_{1\mathcal{U},0\mathcal{U}} = c_t^2\varrho_{12,02}, \quad \rho^{(t)}_{1\mathcal{U},0\mathcal{P}} = -c_t s_t^2\varrho_{10,01}, \\
\rho^{(t)}_{1\mathcal{U},10} &= c_t^2\varrho_{12,10}, \quad \rho^{(t)}_{1\mathcal{U},1\mathcal{D}} = c_t^2\varrho_{12,11}, \quad \rho^{(t)}_{1\mathcal{U},1\mathcal{U}} = c_t^2 s_t^2\varrho_{10,10}, \quad \rho^{(t)}_{1\mathcal{U},1\mathcal{P}} = -c_t^2 s_t\varrho_{10,112}, \\
\rho^{(t)}_{1\mathcal{P},00} &= 0, \quad \rho^{(t)}_{1\mathcal{P},0\mathcal{D}} = c_t^2 s_t^2\varrho_{12,00}, \quad \rho^{(t)}_{1\mathcal{P},0\mathcal{U}} = -c_t s_t^2\varrho_{11,00}, \quad \rho^{(t)}_{1\mathcal{P},0\mathcal{P}} = s_t^2(\varrho_{11,01} + \varrho_{12,02}), \\
\rho^{(t)}_{1\mathcal{P},10} &= 0, \rho^{(t)}_{1\mathcal{P},1\mathcal{D}} = c_t^2\varrho_{12,11}, \quad \rho^{(t)}_{1\mathcal{P},1\mathcal{U}} = -c_t s_t^2\varrho_{11,10}, \quad \rho^{(t)}_{1\mathcal{P},1\mathcal{P}} = s_t^2(\varrho_{11,12} + \varrho_{12,12}),
\end{aligned} \quad (10)$$

   where $c_t = \cos r_t$ and $s_t = sin\ r_t$.

3. *Both subsystems are accelerated:*
   In this case, we assume that both subsystems are accelerated. The final state of the

system in the first region I of the Rindler space is defined by a matrix of size 8×8 elements. The elements of this matrix are given by,

$$\begin{aligned}
\rho^{(qt)}_{00,jk} &= \mathcal{B}^{(qt)}_{\ell_1}, \quad \ell_1 = 1..3, \quad jk = 00, 0\mathcal{D}, 0\mathcal{U}, \quad \rho^{(qt)}_{00,jk} = \mathcal{B}_{\ell_2}, \quad \ell_2 = 4,5, \quad jk = 10, 1\mathcal{D}, \\
\rho^{(qt)}_{00,12} &= \varrho_{00,1\mathcal{P}} = 0, \quad \rho^{(qt)}_{01,jk} = \mathcal{B}_{\ell_3}, \quad \ell_3 = \mathcal{B}_6 ... \mathcal{B}_{13}, \quad jk = 00, 0\mathcal{D}, 0\mathcal{U}, 0\mathcal{P}, 10, 1\mathcal{D}, 1\mathcal{U}, 1\mathcal{P}, \\
\rho^{(qt)}_{02,jk} &= \mathcal{B}_{\ell_4}, \quad \ell_4 = \mathcal{B}_{14} ... \mathcal{B}_{21}, \quad jk = 00, 0\mathcal{D}, 0\mathcal{U}, 0\mathcal{P}, 10, 1\mathcal{D}, 1\mathcal{U}, 1\mathcal{P}, \quad \rho^{(qt)}_{0\mathcal{P},00} = \rho_{0\mathcal{P},10} = 0, \\
\varrho^{(qt)}_{0\mathcal{P},jk} &= \mathcal{B}_{\ell_5}, \quad \ell_5 = \mathcal{B}_{22} ... \mathcal{B}_{27}, \quad jk = 0\mathcal{D}, 0\mathcal{U}, 0\mathcal{P}, 1\mathcal{D}, 1\mathcal{U}, 1\mathcal{P} \quad \rho^{(qt)}_{10,0\mathcal{P}} = \rho^{(qt)}_{10,1\mathcal{P}} = 0, \\
\rho^{(qt)}_{10,jk} &= \mathcal{B}_{\ell_6}, \quad \ell_6 = \mathcal{B}_{28} ... \mathcal{B}_{33}, \quad jk = 00, 0\mathcal{D}, 0\mathcal{U}, 10, 1\mathcal{D}, 1\mathcal{U}, \quad \rho_{11,00} = 0, \\
\rho^{(qt)}_{1\mathcal{D},jk} &= \mathcal{B}_{\ell_7}, \quad \ell_7 = \mathcal{B}_{34} ... \mathcal{B}_{40}, \quad jk = 0\mathcal{D}, 0\mathcal{U}, 0\mathcal{P}, 10, 1\mathcal{D}, 1\mathcal{U}, 1\mathcal{P}, \\
\rho^{(qt)}_{1\mathcal{U},jk} &= \mathcal{B}_{\ell_8}, \quad \ell_8 = \mathcal{B}_{41} ... \mathcal{B}_{48}, \quad jk = 00, 0\mathcal{D}, 0\mathcal{U}, 10, 1\mathcal{D}, 1\mathcal{U}, 1\mathcal{P}, \\
\rho^{(qt)}_{1\mathcal{P},00} &= \rho^{(qt)}_{1\mathcal{P},0\mathcal{U}} = \rho^{(qt)}_{1\mathcal{P},10} = 0, \quad \rho^{(qt)}_{1\mathcal{P},0\mathcal{D}} = \mathcal{B}_{48}, \quad \rho^{(qt)}_{1\mathcal{P},0\mathcal{P}} = \mathcal{B}_{49}, \\
\rho^{(qt)}_{1\mathcal{P},jk} &= \mathcal{B}_{\ell_9}, \quad \ell_9 = 50...52, \quad jk = 1\mathcal{D}, 1\mathcal{U}, 1\mathcal{P}, \qquad (11)
\end{aligned}$$

where $\mathcal{B}_i$ are given in the appendix (B). For example $\rho^{(qt)}_{00,00} = \mathcal{B}^{(qt)}_1$, $\rho^{(qt)}_{10,0\mathcal{D}} = \mathcal{B}^{(qt)}_{29}$, $\rho^{(qt)}_{1\mathcal{P},1\mathcal{P}} = \mathcal{B}^{(qt)}_{52}$ and so on.

3 Dynamics of Entanglement

To quantify the degree of entanglement $\mathcal{E}$ which is contained in the accelerated system, we use the negativity as a measure. For any composite system ρ_{ab} consists of two subsystems with different dimensions. In our case, the system a represents the qubit (2-dimensions) and b refers to the qutrit (3-dimensions). The negativity for this system is defined as,

$$\mathcal{E} = max\left(0, \sum_i \lambda_i\right), \qquad (12)$$

where $\lambda_i, i = 1..6$ are the eigenvalues of the partial transpose of $\rho^{T_a}_{ab}$ [24].

3.1 Example one

To investigate the effect of the accelerated particles on the degree of entanglement, we consider a system of qubit-qutrit is defined by,

$$\begin{aligned}
\vec{s} &= (0,0,s_3), \quad \vec{t} = (0,0,t_3,0,0,0,0,0), \\
&\text{and the non zero elements of the correlation matrix is,} \\
c_{11} &= 1, \quad c_{22} = -1, \quad c_{33} = 1. \qquad (13)
\end{aligned}$$

By using (7), we obtain the final state of the qubit-qutrit system in the first region, I of the Rindler space when only the qubit is accelerated. This state is described by 6×6 elements in the computational basis $00, 01, 02, 10, 11$ and 12. The non- zero elements of this accelerated state in region I are given by,

$$\begin{aligned}
\rho^{(q)}_{00,00} &= c_q^2 \varrho_{00,00}, \quad \rho^{(q)}_{00,11} = c_q \varrho_{00,11}, \quad \rho^{(q)}_{01,01} = c_q^2 \varrho_{01,01}, \quad \rho^{(q)}_{01,10} = c_q \varrho_{01,10} \\
\rho^{(q)}_{02,02} &= c_q^2 \varrho_{02,02}, \quad \rho^{(q)}_{10,01} = c \varrho_{00,01}, \quad \rho^{(q)}_{10,10} = \varrho_{10,10} + s_q^2 \varrho_{00,00}, \\
\rho^{(q)}_{11,00} &= c_q \varrho_{11,00}, \quad \rho^{(q)}_{11,11} = \varrho_{11,11} + s_q^2 \varrho_{01,01}, \quad \rho^{(q)}_{12,12} = \varrho_{12,12} + s_q^2 \varrho_{01,02}. \qquad (14)
\end{aligned}$$



Figure 1: The degree of entanglement $\mathcal{E}$, where we assume that $r_q = r_t = r$ and (a) $s_3 = 1$, $t_3 = 1$, and (b)$s_3 = t_3 = 0$. The solid and the dot lines shows the behavior of entanglement when only the qubit and only the qutrit are accelerated respectively.

Similarly, If only the qutrit is accelerated, then the non zero elements of the final state of the accelerated system in the first region of Rindler space can be written in the basis $|\ell, 0\rangle, |\ell, \mathcal{D}\rangle, |\ell, \mathcal{U}\rangle, |\ell, \mathcal{P}\rangle$, ($\ell = 0, 1$) as,

$$\begin{aligned}
\rho^{(t)}_{00,00} &= c_t^4 \varrho_{00,00}, \quad \rho^{(t)}_{00,1\mathcal{D}} = c_t^3 \varrho_{00,11}, \quad \rho^{(t)}_{0\mathcal{D},0\mathcal{D}} = c_t^2 \varrho_{01,01} + c_t^2 s_t^2, \quad \rho^{(t)}_{0\mathcal{D},10} = c_t^3 \varrho_{01,10}, \\
\rho^{(t)}_{0\mathcal{P},0\mathcal{P}} &= s_t^4 \varrho_{00,00} + s_t^2(\varrho_{01,01} + \varrho_{02,02}), \quad \rho^{(t)}_{0\mathcal{P}p,1\mathcal{U}} = -c_t s_t^2 \varrho_{01,10}, \quad \rho^{(t)}_{02,1p} = -c_t^2 s_t^2 \varrho_{00,11}, \\
\rho^{(t)}_{0\mathcal{U},0\mathcal{U}} &= c_t^2 s_t^2 \varrho_{00,00} + c_t^2 \varrho_{02,02}, \quad \rho^{(t)}_{10,0\mathcal{D}} = c_t^3 \varrho_{10,01}, \quad \rho^{(t)}_{10,10} = c_t^4 \varrho_{10,10}, \\
\rho^{(t)}_{1\mathcal{D},00} &= c_t^2 \varrho_{11,00}, \quad \rho^{(t)}_{1\mathcal{D},1\mathcal{D}} = c_t^2 \varrho_{11,11} + c_t^2 s \varrho_{10,10}, \quad \rho^{(t)}_{1\mathcal{U},1\mathcal{U}} = c_t^2 \varrho_{12,12} + c_t^2 s_t^2 \varrho_{10,10}, \\
\rho^{(t)}_{1\mathcal{U},0\mathcal{P}} &= -c_t s_t^2 \varrho_{10,01}, \quad \rho^{(t)}_{1\mathcal{P},0\mathcal{U}} = -c_t s_t^2 \varrho_{11,00}, \quad \rho^{(t)}_{1\mathcal{P},1\mathcal{P}} = s_t^4 \varrho_{10,10} + s_t^2(\varrho_{11,11} + \varrho_{12,12}).
\end{aligned} \tag{15}$$

Finally, let us assume that both particles are accelerated. In this case, the elements of the final state in the first region I_1 are given by,

$$\begin{aligned}
\rho^{(qt)}_{00,00} &= c_t^4 c_q^2 \varrho_{00,00}, \quad \rho^{(qt)}_{0\mathcal{D},0\mathcal{D}} = c_q^2 c_t^2(\varrho_{01,01} + s_t^2), \quad \rho^{(qt)}_{10,01} = c_q c_t^3(\varrho_{10,01} + \varrho_{02,01}), \\
\rho^{(qt)}_{0\mathcal{U},0\mathcal{U}} &= c_q^2 c_t^2(\varrho_{02,02} + \varrho_{00,00} s_t^2), \quad \rho^{(qt)}_{1\mathcal{P},0\mathcal{U}} = -c_q c_t s_t^2 \varrho_{11,00}, \\
\rho^{(qt)}_{0\mathcal{P},0\mathcal{P}} &= c_q^2 s_t^2(s_q^2 \varrho_{00,00} + \varrho_{01,01} + \varrho_{02,02}), \quad \rho^{(qt)}_{1\mathcal{U},0\mathcal{P}} = -c_q c_t s_t^2 \varrho_{10,01}, \quad \rho_{01,10} = c_q c_t^3 \varrho_{01,10}, \\
\rho^{(qt)}_{10,10} &= c_t^4(s_q^2 \varrho_{00,00} + c_t^4 \varrho_{10,10}), \quad \rho^{(qt)}_{00,1\mathcal{D}} = c_q c_t^3 \varrho_{00,11}, \\
\rho^{(qt)}_{1\mathcal{D},1\mathcal{D}} &= s_q^2 c_t^2(s_q^2 + \varrho_{01,01}) + c_t^2(\varrho_{11,11} + s_t^2 \varrho_{10,10}), \quad \rho^{(qt)}_{0\mathcal{P},1\mathcal{U}} = -c_q c_t s_t^2 \varrho_{01,10}, \\
\rho^{(qt)}_{1\mathcal{U},1\mathcal{U}} &= s_q^2 c_t^2(\varrho_{02,02} + s_t^2 \varrho_{00,00}) + c_t^2(\varrho_{12,12} + s_t^2 \varrho_{10,10}), \quad \rho^{(qt)}_{0\mathcal{U},1\mathcal{P}} = -c_q c_t s_t^2 \varrho_{00,11}, \\
\rho^{(qt)}_{1\mathcal{P},1\mathcal{P}} &= s_q^2 s_t^2(\varrho_{00,00} s^2 + \varrho_{01,01} + \varrho_{02,02}) + s_t^2(s_t^2 \varrho_{10,10} + \varrho_{11,11} + \varrho_{11,01}),
\end{aligned} \tag{16}$$

where the suffix "q", "t" and "qt" refer to the accelerated qubit, qutrit and qubit-qutrit, respectively.

The behavior of entanglement, $\mathcal{E}$ is described in Fig.(1a), where we assume that both subsystems are initially polarized on the third direction, namely we set $s_3 = t_3 = 1$. It is clear that, the entanglement decreases as the acceleration of the accelerated subsystems increase. The decay rate of entanglement depends on the accelerated system. However, if the smaller dimensional system is accelerated, then the decay rate of entanglement is smaller than the decay which is caused by accelerating the larger dimensional system. These results are clearly seen by comparing the behavior of endamagement when only the qubit is accelerated (solid-curve)and its behavior when only the qutrit is accelerated (dash-curve). Moreover, the decay rate of entanglement is larger if both subsystems are accelerated.

In Fig.(1b), we consider another system , where we set $s_2 = t_3 = 0$. The general behavior is similar to that shown in Fig.(1a), namely, the entanglement decreases as the accelerations of the subsystem increase. However, the decay rate of entanglement is larger than that depicted in Fig.(1a). This shows that, the entanglement of the initial system plays an essential roles on the degree of entanglement of the accelerated systems.

3.2 One-Parameter Family

This class of qubit-qutrit system is known by one parameter family. The density operator of this system is given by,

$$\begin{aligned}\rho^{(1)}_{qt} &= |0\rangle_q\langle 0|\Big\{\frac{p}{2}\left(|0\rangle_t\langle 0| + |1\rangle_t\langle 1|\right) + \frac{1-2p}{2}|2\rangle_t\langle 2|\Big\} + |1\rangle_q\langle 0|\Big\{\frac{p}{2}|2\rangle_t\langle 0|\Big\} \\ &+ |0\rangle_q\langle 1|\Big\{\frac{p}{2}|0\rangle_t\langle 2| + \frac{1-2p}{2}|2\rangle_t\langle 0|\Big\} + |1\rangle_q\langle 1|\Big\{\frac{p}{2}|1\rangle_t\langle 1| + \frac{1-2p}{2}|0\rangle_t\langle 0|\Big\}, \quad (17)\end{aligned}$$

where $0 \leq p \leq \frac{1}{2}$ and the subscript "q" refers to the qubit while "t" refers to the qutrit. If we assume that, only the qubit is accelerated, then the density operator of the final state in the first region (I) is defined by the following non-zero elements, of a matrix of size 2×3,

$$\begin{aligned}\rho^{(1_q)}_{00,00} &= \frac{p}{2}c_q^2, \quad \rho^{(1_q)}_{00,12} = \frac{p}{2}c_q, \quad \rho^{(1_q)}_{01,01} = \frac{p}{2}c_q^2, \quad \rho^{(1_q)}_{00,12} = \rho_{12,00}, \\ \rho^{(1_q)}_{02,02} &= \frac{1-2p}{2}c_q^2, \quad \rho^{(1_q)}_{10,02} = \frac{1-2p}{2}c_q, \quad \rho^{(1_q)}_{10,10} = \left(\frac{1-2p}{2} + \frac{p}{2}s_q^2\right), \\ \rho^{(1_q)}_{02,10} &= \rho_{10,02}, \quad \rho^{(1_q)}_{11,11} = \frac{p}{2}(1+s_q^2), \quad \rho^{(1_q)}_{12,12} = \left(\frac{1-2p}{2}s_q^2 + \frac{p}{2}\right). \quad (18)\end{aligned}$$

On the other hand, if only the qutrit is accelerated, then the final state of the total system in the first region is defined by a matrix of size 8×8. The non-zero elements are given by,

$$\begin{aligned}\rho^{(1_t)}_{00,00} &= \frac{p}{2}c_t^4, \quad \rho^{(1_t)}_{12,00} = \frac{p}{2}c_t^3, \quad \rho^{(1_t)}_{0\mathcal{D},0\mathcal{D}} = \frac{p}{2}c_t^2(1+s_t^2), \quad \rho^{(1_t)}_{1\mathcal{P},0\mathcal{D}} = \frac{p}{2}c_t s_t^2, \\ \rho^{(1_t)}_{0\mathcal{U},0\mathcal{U}} &= c_t^2\left(\frac{1-2p}{2} + \frac{p}{2}s_t^2\right), \quad \rho^{(1_t)}_{10,02} = \frac{1-2p}{2}c_t^3, \quad \rho^{(1_t)}_{0\mathcal{P},0\mathcal{P}} = s_t^2\left(\frac{p}{2}(1+s_t^2) + \frac{1-2p}{2}\right), \\ \rho^{(1_t)}_{1\mathcal{D},0\mathcal{P}} &= \frac{1-2p}{2}c_t s_t^2, \quad \rho^{(1_t)}_{10,10} = \frac{1-2p}{2}c_t^4, \quad \rho^{(1_t)}_{0\mathcal{U},10} = \rho^{(1_t)}_{10,0\mathcal{U}}, \quad \rho^{(1_t)}_{1\mathcal{D},1\mathcal{D}} = \frac{1-2p}{2}c_t^2 s_t^2, \\ \rho^{(1_t)}_{0\mathcal{P},1\mathcal{D}} &= \frac{1-2p}{2}c_t s_t^2, \quad \rho^{(1_t)}_{00,1\mathcal{U}} = \rho^{(1_t)}_{12,00}, \quad \rho^{(1_t)}_{1\mathcal{U},1\mathcal{U}} = c_t^2\left(\frac{p}{2} + \frac{1-2p}{p}s_t^2\right), \\ \rho^{(1_t)}_{0\mathcal{D},1\mathcal{P}} &= \frac{p}{2}c_t s_t^2, \quad \rho^{(1_t)}_{1\mathcal{P},1\mathcal{P}} = s_t^2\left(p + \frac{1-2p}{2}s_t^2\right). \quad (19)\end{aligned}$$



Figure 2: The entanglement of one parameter family evaluated at $p = 0.5, r_q = r_t = r$. The solid, dash and dot curves represent the entanglement $\mathcal{E}$, when only the qubit , only the qutrit, and both of them are accelerated.

Finally, if both subsystems are accelerated, then the final state in the first region, I is defined by the following non-zero elements of a matrix of 8×8,

$$\begin{aligned}
\rho^{(1_{qt})}_{00,00} &= c_q^2 \rho^{(1_t)}_{00,00}, \quad \rho^{(1_{qt})}_{1\mathcal{D},00} = \frac{p}{2} c_q c_t^3, \quad \rho^{(1_{qt})}_{0\mathcal{D},0\mathcal{D}} = c_q^2 \rho^{(1_t)}_{0\mathcal{D},0\mathcal{D}}, \quad \rho^{(1_{qt})}_{1\mathcal{P},0\mathcal{D}} = c_q \rho^{(1_t)}_{1\mathcal{P},0\mathcal{D}}, \\
\rho^{(1_{qt})}_{0\mathcal{U},0\mathcal{U}} &= c_q^2 \rho^{(1_t)}_{0\mathcal{U},0\mathcal{U}}, \quad \rho^{(1_{qt})}_{10,0\mathcal{U}} = c_q \rho^{(1_t)}_{10,0\mathcal{U}}, \quad \rho^{(1_{qt})}_{0\mathcal{P},0\mathcal{P}} = c_q^2 \rho^{(1_t)}_{0\mathcal{P},0\mathcal{P}}, \quad \rho^{(1_{qt})}_{1\mathcal{D},0\mathcal{P}} = c_q \rho^{(1_t)}_{1\mathcal{D},0\mathcal{P}}, \\
\rho^{(1_{qt})}_{10,10} &= \rho^{(1_t)}_{10,10} + \frac{p}{2} c_t^4 s_q^2, \quad \rho^{(1_{qt})}_{0\mathcal{D},10} = c_q \rho^{(1_t)}_{10,0\mathcal{U}}, \quad \rho^{(1_{qt})}_{1\mathcal{D},1\mathcal{D}} = \rho^{(1_t)}_{1\mathcal{D},1\mathcal{D}} + c_t^2 \frac{p}{2} \left(1 + s_q^2 + s_q^2 s_t^2\right), \\
\rho^{(1_{qt})}_{0\mathcal{P},1\mathcal{D}} &= c_q \rho^{(1_t)}_{0\mathcal{P},1\mathcal{D}}, \quad \rho^{(1_{qt})}_{1\mathcal{U},1\mathcal{U}} = \rho^{(1_t)}_{1\mathcal{U},1\mathcal{U}} + c_t^2 \left(\frac{p}{2}(1 + s_t^2 s_q^2) + \frac{1-2p}{2} s_q^2\right), \quad \rho^{(1_{qt})}_{00,1\mathcal{U}} = c_q \rho^{(1_t)}_{00,1\mathcal{U}}, \\
\rho^{(1_{qt})}_{1\mathcal{P},1\mathcal{P}} &= \rho^{(1_t)}_{1\mathcal{P},1\mathcal{P}} + s_q^2 s_t^2 \left(\frac{p}{2} s_t^2 + \frac{1-p}{2}\right), \quad \rho^{(1_{qt})}_{01,1\mathcal{P}} = c_q \rho^{(1_t)}_{0\mathcal{D},1\mathcal{P}}. \qquad (20)
\end{aligned}$$

The behavior of entanglement for this case is shown in Fig.(2), where we initially start with a system defined by $p = 0.5$. The general behavior is similar to that shown in Fig.(1), namely the entanglement decays as the the acceleration of the accelerated subsystem increases. The upper and lower bounds of entanglement depend on the accelerated subsystem and the initial degree of entanglement. However, if the qubit is allowed to be accelerated, then the upper bounds of entanglement are always larger than that shown if one allows the qutrit to be accelerated. The decay rate of entanglement increases if both particles are accelerated, where we considered both particles are accelerated with the same acceleration (i.e., $r_q = r_t = r$).

3.3 Two-Parameters family

In this subsection we consider the second example which is known by a two-parameter family. In the computational basis, this class can be written as,

$$\begin{aligned}\rho^{(2)}_{qt} &= |0\rangle_q\langle 0|\Big\{\beta|0\rangle_t\langle 0| + \frac{\beta+\gamma}{2}|1\rangle_t\langle 1| + \alpha|2\rangle_t\langle 2|\Big\} + |1\rangle_q\langle 0|\Big\{\frac{\beta-\gamma}{2}|0\rangle_t\langle 1|\Big\} \\ &\quad + |0\rangle_q\langle 1|\Big\{\frac{\beta-\gamma}{2}|1\rangle_t\langle 0|\Big\} + |1\rangle_q\langle 1|\Big\{\frac{\beta+\gamma}{2}|0\rangle_t\langle 0| + \beta|1\rangle\langle 1| + \alpha|2\rangle_t\langle 2|\Big\}, \qquad (21)\end{aligned}$$

where $\gamma + 2\alpha + 3\beta = 1$ and the suffix 2 refers to the two-parameter. If it assumed that only the qubit is accelerated, then the final state in the first region, I of the Rindler space is described by the following non-zero-elements of a matrix of size 6×6,

$$\begin{aligned}\rho^{(2_q)}_{00,00} &= \beta c_q^2, \quad \rho^{(2_q)}_{01,01} = \frac{\beta+\gamma}{2}c_q^2, \quad \rho^{(2_q)}_{10,01} = \frac{\beta-\gamma}{2}c_q, \\ \rho^{(2_q)}_{02,02} &= \alpha c_q^2, \quad \rho^{(2_q)}_{01,10} = \frac{\beta-\gamma}{2}c_q, \quad \rho_{10,10} = \beta s_q^2, \\ \rho^{(2_q)}_{11,11} &= \left(\beta + \frac{\beta+\gamma}{2}s_q^2\right), \quad \rho^{(2_q)}_{12,12} = \alpha(1+s_q^2). \qquad (22)\end{aligned}$$

On the other hand, if we consider only the qutrit is accelerated then, the final state in the first Rindler region is defined by the following non-zero elements,

$$\begin{aligned}\rho^{(2_t)}_{00,00} &= \beta c_t^4, \quad \rho^{(2_t)}_{01,0\mathcal{D}} = c_t^2\left(\beta s_t^2 + \frac{\beta+\gamma}{2}\right), \quad \rho^{(2_t)}_{10,0\mathcal{D}} = \frac{\beta-\gamma}{2}c_t^3, \\ \rho^{(2_t)}_{0\mathcal{U},0\mathcal{U}} &= c_t^2(\alpha + \beta s_t^2), \quad \rho^{(2_t)}_{0\mathcal{P},0\mathcal{P}} = s_t^2(\left(\alpha + \frac{3\beta+\gamma}{2}s_t^2\right), \quad \rho^{(2_t)}_{1\mathcal{U},0\mathcal{P}} = -\frac{\beta-\gamma}{2}c_t s_t^2, \\ \rho^{(2_t)}_{0\mathcal{D},10} &= \frac{\beta-\gamma}{2}c_t^3, \quad \rho^{(2_t)}_{10,10} = \frac{\beta+\gamma}{2}c_t^4, \quad \rho^{(2_t)}_{1\mathcal{D},1\mathcal{D}} = c_t^2\left(\beta + \frac{\beta+\gamma}{2}s_t^2\right), \quad \rho^{(2_t)}_{0\mathcal{P},1\mathcal{U}} = \rho^{(2_t)}_{1\mathcal{U},0\mathcal{P}} \\ \rho^{(2_t)}_{1\mathcal{U},1\mathcal{U}} &= c_t^2\left(\alpha + \frac{\beta+\gamma}{2}s_t^2\right), \quad \rho^{(2_t)}_{1\mathcal{P},1\mathcal{P}} = s_t^2\left(\alpha + \frac{3\beta+\gamma}{2}s_t^2\right). \qquad (23)\end{aligned}$$

Finally, when both subsystems are accelerated, then the final state in the first region, I is defined by a matrix of size 8×8 elements. These elements can be written by (22) as,

$$\begin{aligned}\rho^{(2_{qt})}_{00,00} &= c_q^2\rho^{(2_t)}_{00,00}, \quad \rho^{(2_{qt})}_{0\mathcal{D},0\mathcal{D}} = c_q^2\rho^{(2_t)}_{0\mathcal{D},0\mathcal{D}}, \quad \rho^{(2_{qt})}_{10,0\mathcal{D}} = c_q\rho^{(2_t)}_{10,0\mathcal{D}}, \quad \rho^{(2_{qt})}_{0\mathcal{U},0\mathcal{U}} = c_q^2\rho^{(2_t)}_{0\mathcal{U},0\mathcal{U}} \\ \rho^{(2_{qt})}_{0\mathcal{P},0\mathcal{P}} &= c_q^2\rho^{(2_t)}_{0\mathcal{P},0\mathcal{P}}, \quad \rho^{(2_{qt})}_{1\mathcal{U},0\mathcal{P}} = c_q\rho^{(2_t)}_{1\mathcal{U},0\mathcal{P}}, \quad \rho^{(2_{qt})}_{0\mathcal{D}10} = c_q\rho^{(2_t)}_{0\mathcal{D},10}, \quad \rho^{(2_{qt})}_{10,10} = s_q^2\rho^{(2_t)}_{10,10}, \\ \rho^{(2_{qt})}_{1\mathcal{D},1\mathcal{D}} &= \rho^{(2_t)}_{11,11} + s_q^2\rho^{(2_t)}_{0\mathcal{D},0\mathcal{D}}, \quad \rho^{(2_{qt})}_{0\mathcal{P},1\mathcal{U}} = \rho^{(2_{qt})}_{1\mathcal{U},0\mathcal{P}}, \quad \rho^{(2_{qt})}_{1\mathcal{U},1\mathcal{U}} = \rho^{(2_t)}_{1\mathcal{U},1\mathcal{U}} + s_q^2\rho^{(2_t)}_{0\mathcal{U},0\mathcal{U}}, \\ \rho^{(2_{qt})}_{1p,1p} &= \rho^{(2_t)}_{1\mathcal{P},1\mathcal{P}} + s_q^2\rho^{(2_t)}_{0\mathcal{P},0\mathcal{P}}, \qquad (24)\end{aligned}$$

The behavior of entanglement for this class is described in Fig.(3), where all the three possibilities are considered. It is assumed that, the system is initially prepared in maximum entangled state $\alpha = \beta = 0.5$. Therefore the entanglement is maximum i.e., $\mathcal{E} = 1$ at $r_q = r_t = 0$. It is clear that, the behavior of entanglement is similar to that predicated for one parameter family. However, the entanglement decays as the acceleration of any subsystem increases, where we consider that, both of the subsystems have the same acceleration $(r_q = r_t)$. The rate of entanglement decay increases as the dimensions of the accelerated subsystems increase. Moreover, if both subsystems are accelerated, then the entanglement decreases faster.



Figure 3: The dynamics of entanglement in the first region for qubit-qutrit system of two-parameter family with $\alpha = \beta = 0.5$. The solid, dash and dot-curves represent $\mathcal{E}$, when only the qubit, qutrit and both qubit and qutrit are accelerated.

4 Conclusion

In this contribution, we consider a general system composite of two different dimensions. One of them is a two-dimensional qubit and the other is of three dimensional qutrit. It is assumed that, one or both subsystems are accelerated simultaneously or non- simultaneously. The density operator of the final state in the first region of the accelerated is obtained analytical in a general form. Different examples are introduced to show the main task of this manuscript.

Our results show that, the entanglement decays as the acceleration of the accelerated particle increases. However, the decay rate depends on the dimensions of the accelerated subsystem. If only the qubit is accelerated, then the entanglement decays gradually to reach its non-zero values at infinity. On the other hand, if we allow only the qutrit to be accelerated, then the entanglement decays faster and its upper bounds always smaller than that depicted for accelerating qubit. Moreover, if one allows for both subsystems to be accelerated, then the decay rate of entanglement is larger.

The idea is shown explicitly by considering three different and common examples. In the first example, we assume that both subsystems are polarized in $z-$axis. It is clear that, the decay of entanglement depends on the initial correlation between the two subsystems. The second and third examples are known by one and two-parameters families. These examples have been investigated widely in the context of quantum information. It is shown that the decay rate of entanglement depends on the initial degree of entanglement in addition to the dimensions of the accelerated subsystem.

In conclusion: the decay of entanglement depends on the dimensions of the accelerated subsystems, where the decay rate of entanglement increases when the larger dimensional subsystem is accelerated. The robustness of the accelerated system depends on the initial quantum correlation between the two subsystems. We expect that these results could be important in building an accelerated quantum network consists of different dimensions nodes.

Appendices

A The elements of the initial state

The elements of the initial system (1) in the basis $|00\rangle, |01\rangle, |02\rangle, |10\rangle, |11\rangle$ and $|12\rangle$ are given by.

$$\begin{aligned}
\mathcal{A}_1 &= (1+s_3+t_3-c_{33}+t_8/\sqrt{3}-c_{38}/\sqrt{3})/6, \quad \mathcal{A}_2=(t_1-it_2-c_{31}+ic_{32})/6,\\
\mathcal{A}_3 &= (t_4-it_5-c_{34}+ic_{35})/6, \quad \mathcal{A}_4=(s_1+is_2+c_{13}+ic_{23}+c_{18}/\sqrt{3}+ic_{28}/\sqrt{3})/6,\\
\mathcal{A}_5 &= (c_{11}-c_{22}+ic_{21}-ic_{12})/6, \quad \mathcal{A}_6=(c_{14}+c_{25}+ic_{24}-ic_{15})/6,\\
\mathcal{A}_7 &= (t_1+it_2-c_{31}-ic_{32})/6, \quad \mathcal{A}_8=(1-s_3-t_3+t_8/\sqrt{3}+c_{33}-c_{38}/\sqrt{3})/6,\\
\mathcal{A}_9 &= (t_6-it_7-c_{36}+ic_{37})/6, \quad \mathcal{A}_{10}=(c_{11}-c_{22}+ic_{12}-ic_{21})/6,\\
\mathcal{A}_{11} &= (s_1+is_2-c_{13}+-ic_{23}+c_{18}/\sqrt{3}+ic_{28}/\sqrt{3})/6, \quad \mathcal{A}_{12}=(c_{16}+c_{27}+ic_{26}-ic_{17})/6,\\
\mathcal{A}_{13} &= (t_4+it_5-c_{34}-ic_{35})/6, \quad \mathcal{A}_{14}=(t_6+it_7-c_{36}-ic_{37})/6,\\
\mathcal{A}_{15} &= (1-s_3-2t_8/\sqrt{3}+2c_{38}/\sqrt{3})/6, \quad \mathcal{A}_{16}=(c_{14}-c_{25}+ic_{15}+ic_{24})/6,\\
\mathcal{A}_{17} &= (c_{16}-c_{27}+ic_{17}+ic_{26})/6, \quad \mathcal{A}_{18}=(s_1+is_2-2c_{28}/\sqrt{3}-2c_{18}/\sqrt{3})/6,\\
\mathcal{A}_{19} &= (s_1+is_2+c_{13}-ic_{23}+c_{18}/\sqrt{3}-ic_{28}/\sqrt{3})/6, \quad \mathcal{A}_{20}=(c_{11}-c_{22}-ic_{12}-ic_{21})/6,\\
\mathcal{A}_{21} &= (c_{14}-c_{25}-ic_{15}-ic_{24})/6, \quad \mathcal{A}_{22}=(1+s_3+t_3+t_8/\sqrt{3}+c_{33}+c_{38}/\sqrt{3})/6,\\
\mathcal{A}_{23} &= (t_1-it_2+c_{31}-ic_{32})/6, \quad \mathcal{A}_{24}=(t_4-it_5+c_{34}-ic_{35})/6,\\
\mathcal{A}_{25} &= (c_{11}+c_{22}+ic_{12}-ic_{21})/6, \quad \mathcal{A}_{26}=(s_1-is_2-c_{13}+ic_{23}+c_{18}/\sqrt{3}-ic_{28}/\sqrt{3})/6,\\
\mathcal{A}_{27} &= (c_{16}-c_{27}-ic_{17}-ic_{26})/6, \quad \mathcal{A}_{28}=-(t_1+it_2+c_{31}+ic_{32})/6,\\
\mathcal{A}_{29} &= (1+s_3-t_3+t_8/\sqrt{3}-c_{33}+c_{38}/\sqrt{3})/6, \quad \mathcal{A}_{30}=(t_6-it_7+c_{36}-ic_{37})/6,\\
\mathcal{A}_{31} &= (c_{14}+c_{25}+ic_{15}+ic_{24})/6, \quad \mathcal{A}_{32}=(c_{16}+c_{27}+ic_{17}-ic_{26})/6,\\
\mathcal{A}_{33} &= (s_1-is_2+2ic_{28}/\sqrt{3}-2c_{18}/\sqrt{3})/6, \quad \mathcal{A}_{34}=(t_6+it_7+c_{36}+ic_{37})/6,\\
\mathcal{A}_{35} &= (t_4+it_5+c_{34}+ic_{35})/6, \quad \mathcal{A}_{36}=(1+s_3-2t_8/\sqrt{3}-2c_{38}/\sqrt{3})/6. \qquad (25)
\end{aligned}$$

B Both subsystems are accelerated

The parameters $\mathcal{B}_i, i, 1..52$ which are appeared in Eq.(11) are give explicitly as,

$$\begin{aligned}
\mathcal{B}_1 &= c_q^2c_t^4\mathcal{A}_1, \quad \mathcal{B}_2=c_q^2\mathcal{A}_2, \quad \mathcal{B}_3=c_q^2\mathcal{A}_3+c_q\mathcal{A}_6, \quad \mathcal{B}_4=c_qc_t^4\mathcal{A}_4, \quad \mathcal{B}_5=c_qc_t^3\mathcal{A}_5,\\
\mathcal{B}_6 &= c_q^2c_t^3\mathcal{A}_6, \quad \mathcal{B}_7=c_q^2c_t^2(\mathcal{A}_8+s_t^2\mathcal{A}_1), \quad \mathcal{B}_8=c^qc_t^2\mathcal{A}_9, \quad \mathcal{B}_9=c_q^2c_ts_t^2\mathcal{A}_3, \quad \mathcal{B}_{10}=c_qc_t^3\mathcal{A}_{10}\\
\mathcal{B}_{11} &= c_qc_t^2(\mathcal{A}_{11}+s_t^2\mathcal{A}_4), \quad \mathcal{B}_{12}=c_qc_t^2\mathcal{A}_{12}, \quad \mathcal{B}_{13}=c_qs_t^2\mathcal{A}_6, \quad \mathcal{B}_{14}=c_q^2c_t^3\mathcal{A}_{13}, \quad \mathcal{B}_{15}=c_q^2c_t^2\mathcal{A}_{14}\\
\mathcal{B}_{16} &= c^qc_t^2(\mathcal{A}_{15}+s_t^2\mathcal{A}_1), \quad \mathcal{B}_{17}=-c_q^2c_ts_t^2\mathcal{A}_2, \quad \mathcal{B}_{18}=c_qc_t^3\mathcal{A}_{16}, \quad \mathcal{B}_{19}=c_qc_t^2\mathcal{A}_{19},\\
\mathcal{B}_{20} &= c_q(\mathcal{A}_{18}+c_t^2s_t^2\mathcal{A}_4), \quad \mathcal{B}_{21}=-c_qc_t^2s_t^2\mathcal{A}_5, \quad \mathcal{B}_{22}=c_q^2c_ts_t^2\mathcal{A}_{13}, \quad \mathcal{B}_{23}=-c_q^2c_ts_t^2\mathcal{A}_7,\\
\mathcal{B}_{24} &= c_q^2(s_q^2\mathcal{A}_{15}+s_t^2\mathcal{A}_8+s_t^4\mathcal{A}_1), \quad \mathcal{B}_{25}=c_qc_ts_t^2\mathcal{A}_{16}, \quad \mathcal{B}_{26}=-c_qc_ts_t^2\mathcal{A}_{10},\\
\mathcal{B}_{27} &= c_q(s_t^4\mathcal{A}_1+s_t^2\mathcal{A}_{18}+s_t^2\mathcal{A}_{11}), \quad \mathcal{B}_{28}=c_qc_t^3(c_t\mathcal{A}_{19}+\mathcal{A}_{25}), \quad \mathcal{B}_{29}=c_qc_t^3\mathcal{A}_{20}, \quad \mathcal{B}_{30}=c_qc_t^3\mathcal{A}_{21}\\
\mathcal{B}_{31} &= c_t^4(s_q^2\mathcal{A}_1+\mathcal{A}_{22}), \quad \mathcal{B}_{32}=c_t^3(s_q^2\mathcal{A}_2+\mathcal{A}_{23}), \quad \mathcal{B}_{33}=c_t^3(s_q^2\mathcal{A}_3+\mathcal{A}_{24}),\\
\mathcal{B}_{34} &= c_qc_t^2(s_q^2\mathcal{A}_{19}+\mathcal{A}_{26}), \quad \mathcal{B}_{35}=c_qc_t^2\mathcal{A}_{27}, \quad \mathcal{B}_{36}=c_qc_ts_t^2\mathcal{A}_{21}, \quad \mathcal{B}_{37}=c_t^3(s_q^2\mathcal{A}_7+\mathcal{A}_{28}),\\
\mathcal{B}_{38} &= c_q^2s_t^2\mathcal{A}_{22}+c_t^2\mathcal{A}_{29}+s_q^2c_t^2\mathcal{A}_8+s_q^2c_t^2s_t^2\mathcal{A}_1. \quad \mathcal{B}_{39}=c_t^2(s_q^2\mathcal{A}_9+\mathcal{A}_{30}), \quad \mathcal{B}_{40}=c_ts_t^2(s_q^2\mathcal{A}_3+\mathcal{A}_{24}),\\
\mathcal{B}_{41} &= c_qc_t^3\mathcal{A}_{31}, \quad \mathcal{B}_{42}=c_qc_t^2\mathcal{A}_{32}, \quad \mathcal{B}_{43}=c_qc_t^2(s_t^2\mathcal{A}_{19}+\mathcal{A}_{33}), \quad \mathcal{B}_{44}=-c_qc_ts_t^2\mathcal{A}_{20},
\end{aligned}$$

$$\begin{aligned}
\mathcal{B}_{45} &= c_t^3(s_q^2\dashv_{13}+\mathcal{A}_{34}), \quad \mathcal{B}_{46}=c_t(c_t\mathcal{A}_{35}+s_t^2\mathcal{A}_{14}), \quad \mathcal{B}_{47}=c^2s^2\mathcal{A}_{22}+c^2\mathcal{A}_{36}+s_q^2c_t^2\mathcal{A}_{15}+s_q^2c_t^2s_t^2\mathcal{A}_1\\
\mathcal{B}_{48} &= c_qc_ts_q^2(\mathcal{A}_{31}-\mathcal{A}_{25}), \quad \mathcal{B}_{49}=c_qs_t^2(s^2\mathcal{A}_{19}+s_q^2\mathcal{A}_{26}+\mathcal{A}_{33}), \quad \mathcal{B}_{50}=c_ts_t^2(s^2q\mathcal{A}_{13}+\mathcal{A}_{34}),\\
\mathcal{B}_{51} &= -c_ts_t^2(s_q^2\mathcal{A}_7+\mathcal{A}_{28}), \quad \mathcal{B}_{52}=s^2\mathcal{A}_{29}+s_q^2s_t^2(\mathcal{A}_8+\mathcal{A}_{15})+s_q^2s_t^4\mathcal{A}_1+s_t^3\mathcal{A}_{22}+s^2\mathcal{A}_{36}.
\end{aligned} \tag{26}$$

References

[1] M. A. Nielsen, I. L. Chuang" Quantum Computation and Quantum Information", (Cambradge University Press), Cambridge (2000)

[2] H. Eleuch and R. Bennaceur, J. Opt. A: Pure Appl. Opt. **5** 528 (2003); P. K. Jha, H. Eleuch and Y. V. Rostovtsev, Phys. Rev. A **82** 045805 (2010); H. Eleuch, Eur. Phys. J. D **48** 139 (2008).

[3] Eyob A. Sete and H. Eleuch, Phys. Rev. A **82** 043810 (2010) ; H. Jabri, H. Eleuch, T. Djerad, Laser Phys. Lett. **2** 253 (2005) ; K. Berrada, A. Chafik, H. Eleuch and Y. Hassouni Quantum Inf. Process. **9** 13 (2010). ; H. Eleuch, Int. J. Mod. Phys. B **24** 5653 (2010).

[4] E. A. Sete, H. Eleuch and C. H. R. Ooi, J. Opt. Soc. Am. B **31** 2821 (2014). ; K. Berrada, M. El Baz, H. Eleuch and Y. Hassouni, Int. J Mod. Phys. C **21** 291 (2010).

[5] Hai-Rui Wei, Bao-Cang Ren, Tao Li, Ming Hua, Fu-Guo Deng," Commun. Theor. Phys. **57** 983-990 (2012).

[6] G. Karpat and Z. Gedik, Phys. Lett. A **375** 4166-4171 (2011).

[7] J.-Liang Guo, H. Lu and G. Lu Long, Quantum Inf. Process **12** 3421-3435 (2013).

[8] X. Xiao, Phys. Scr. 065102 (7pp) (2014).

[9] P. Alsing, I. F.-Schuller, R. Mann and T. Tessier, Phys. Rev. A **74** 032326 (2006).

[10] J. Wang, J. Deng and J. Jing, phys. Rev. A**81** 052120 (2010).

[11] N. Metwally, J. Mod. Phys. B **27** 1350155 (18pages) (2013).

[12] N. Metwally and A. Sagheer, Quantum Inf. Process **13** 771-780 (2014).

[13] P. Alsing and G. Milburn, Phys. Rev. Lett. **91** 180404 (2003).

[14] P. Alsing D. McMahon and G. Milburn, J. Opt. B:Quantum Seemiclass Opt. **6** S834-S843 (2004).

[15] N. Metwally, J. Opt. Soc. Am B **30** 233-237 (2013).

[16] S. Khan and M. K. Khan, J. Mod. Opt. **58** 918-923 (2011).

[17] Y. Hao, W. Lian-Fu, Commum. Theor. Phys. **59** 17- 21 (2013).

[18] N. Metwally, Phys. Lett. A **375** 4268-4273 (2011).

[19] E. M.-Martinez, I. Fuentes, Phys. Rev. A **83**, 052306 (2011).

[20] D. F. Walls and G. J. Milburn,”Quantum Optics”, Springer-Verlag, N. Y., (1994).

[21] D. E. Bruschi, J. Louko, E. Martn-Martnez, A. Dragan, and I. Fuentes, Phys. Rev. A **82** 042332 (2010).

[22] J. Doukas, E. G. Brown, A. J. Dragan, and R. B. Mann, Phys. Rev. A **87**, 012306 (2013);

[23] J. Leòn and E. M.-Martìnez, Phys. Rev. A **80** 012314 (2009).

[24] G. Karpat and Z. Gedik, Phys. Lett. A **375** 4166-4171 (2011).